Government of Saskatchewan

Crop Report

Published by the Ministry of Agriculture
ISSN 0701 7085
Report number 9, July 3, 2014

For the Period June 24 to 30, 2014

According to Saskatchewan Agriculture's weekly Crop Report, crop development continues to be delayed due to cool and wet weather conditions. Forty-nine per cent of the fall cereals, 61 per cent of the spring cereals, 66 per cent of the oilseeds and 55 per cent of the pulse crops are behind their normal developmental stages.

One Year Ago

Livestock producers had six per cent of the hay crop cut and 91 per cent was in good to excellent condition. Crop conditions were fair to excellent.

Follow the 2014 Crop Report on Twitter @SKAgriculture

Significant amounts of rain fell over much of the province this weekend with some southeastern and east-central areas receiving well over six inches of rain in a few days. Crops in the hardest hit areas have been significantly impacted by localized flooding and saturated field conditions.

Provincial topsoil moisture conditions on cropland are rated as 48 per cent surplus and 52 per cent adequate. Hay land and pasture moisture conditions are rated as 36 per cent surplus, 62 per cent adequate and two per cent short. Warm and dry weather is needed to help alleviate excess moisture.

The majority of crops outside the areas affected by flooding are in fair to excellent condition. Excess moisture, localized flooding, wind and diseases such as root rot and leaf spots have caused the majority of crop damage this week.

Saskatchewan livestock producers have less than one per cent of the 2014 hay crop cut and 98 per cent is rated as fair to excellent in quality. Much of the hay is behind its normal developmental stages for this time of year.

Farmers are busy controlling weeds and beginning haying operations.

SK Crop Development - June 30

	% Ahead	% Normal	% Behind
Fall Cereals	4	47	49
Spring Cereals	1	38	61
Oilseeds	1	33	66
Pulse Crops	1	44	55

For further information, contact Shannon Friesen, PAg,
Cropping Management Specialist, Moose Jaw, Regional Services Branch,
Toll Free: 1-866-457-2377 or 306-694-3592, E-mail: cropreport@gov.sk.ca.
Also available on the Ministry of Agriculture website at www.agriculture.gov.sk.ca.

Agriculture and Agri-Food Canada | Agriculture et Agroalimentaire Canada

SCIC
SASKATCHEWAN CROP INSURANCE CORPORATION

Southeastern Saskatchewan (Crop District 1 – Carnduff, Estevan, Redvers, Moosomin and Kipling areas; Crop District 2 – Weyburn, Milestone, Moose Jaw, Regina and Qu'Appelle areas; Crop District 3ASE – Radville and Lake Alma areas)

SE SK Crop Development - June 30			
	% Ahead	% Normal	% Behind
Fall Cereals	7	34	59
Spring Cereals	0	19	81
Oilseeds	0	20	80
Pulse Crops	0	18	82

Heavy rainfall over the weekend has flooded much of the southeast. There are reports of washed out highways, grid roads, yards and many homes have flooded basements. Access to good roads is minimal for many in the area. Local slough, creek and lake levels have risen significantly. There are reports of stranded livestock in some areas due to flooding. Some areas around Moosomin are reporting as much as 260 mm in just a few days with many other areas reporting amounts exceeding 150 mm. Moosomin has recorded the greatest amount of rainfall since April 1 with 520 mm, the highest for both the region and the province.

Cropland topsoil moisture conditions are rapidly deteriorating and are now rated as 68 per cent surplus and 32 per cent adequate while hay land and pasture topsoil moisture is rated as 63 per cent surplus and 37 per cent adequate. The majority of crops are in saturated fields with many low spots under water. There are reports that some of the water is receding in some areas but warm and dry weather is needed to help the fields dry and for crops to grow efficiently. Crop loss due to excess moisture is likely to occur in many areas, particularly in low spots and field areas that were saturated before the heavy rain.

Prior to the rainfall, crops were in fair to excellent condition, although the majority was behind in development due to excess moisture and cool weather conditions. For those producers who are still able to get onto their land, in-crop spraying continues as weather and crop staging allow. Besides the excess moisture and localized flooding damage this week, other sources of crop damage include cutworms and root rots. There are also reports of alfalfa weevils in some stands. Haying is just beginning and 89 per cent of the hay crop is rated as good in quality at this time.

Farmers are busy cleaning up flooded yards and homes, controlling weeds and getting ready for haying.

Southwestern Saskatchewan (Crop District 3ASW – Coronach, Assiniboia and Ogema areas; Crop District 3AN – Gravelbourg, Mossbank, Mortlach and Central Butte areas; Crop District 3B – Kyle, Swift Current , Shaunavon and Ponteix areas; Crop District 4 – Consul, Maple Creek and Leader areas)

Haying operations are underway in the region with less than one per cent cut; 88 per cent is rated as good quality at this time. Much of the region received rainfall this week which has further delayed fieldwork. For some producers the rain was welcomed while others' fields are now too wet for in-crop spraying to continue. There are reports that aerial herbicide applications are increasing in many areas.

SW SK Crop Development - June 30			
	% Ahead	% Normal	% Behind
Fall Cereals	0	73	27
Spring Cereals	0	61	39
Oilseeds	0	57	43
Pulse Crops	0	56	44

The Stewart Valley received 57 mm of rain this week. The Cadillac area has received the greatest amount of rain since April 1 (366 mm). Cropland topsoil moisture conditions are rated as 21 per cent surplus, 78 per cent adequate and one per cent short. Hay land and pasture topsoil moisture is rated as nine per cent surplus, 85 per cent adequate and six per cent short. Much of the cropland, hay land and pasture remain short of topsoil moisture in CD 4A.

The majority of crops are at their normal stages of development for this time of year and are in fair to excellent condition. Sun and warm weather is needed to help advance crops and dry excess moisture. Some producers are spraying fungicides on earlier seeded crops. Crop damage this week is attributed to localized flooding, drought, hail and diseases such as root rot.

Farmers are busy controlling weeds, beginning haying operations and hauling grain.

East-Central Saskatchewan (Crop District 5 – Melville, Yorkton, Cupar, Kamsack, Foam Lake, Preeceville and Kelvington areas; Crop District 6A – Lumsden, Craik, Watrous and Clavet areas)

The east-central region received large amounts of rain over the weekend which flooded many fields, yards and homes. There are reports of washed out highways and grid roads; and many homes have flooded basements. Water levels in sloughs, creeks and lakes have risen and damaged surrounding areas.

EC SK Crop Development - June 30			
	% Ahead	% Normal	% Behind
Fall Cereals	1	33	66
Spring Cereals	0	25	75
Oilseeds	1	27	72
Pulse Crops	0	48	52

There are also reports of stranded livestock in some areas due to flooding. The Rhein area reported 191 mm of rain over the course of a few days, bringing its total amount of rain received since April 1 to 477 mm.

Topsoil moisture conditions continue to deteriorate and are now rated as 82 per cent surplus and 18 per cent adequate on cropland. Hay land and pasture topsoil moisture is rated as 73 per cent surplus and 27 per cent adequate. CD 5A is reporting that 89 per cent of the cropland and 78 per cent of the hay land and pasture has surplus topsoil moisture at this time. Much of the crop is in standing water or in fields saturated with excess moisture. Water is receding in some areas, although many fields will not fully recover from the excess moisture stress. Warm and dry weather is now needed to help dry fields and allow for less-impacted crops to develop. Crop loss due to excess moisture is likely to occur in many areas, particularly in low spots and field areas that were saturated prior to the recent rain.

The majority of crops were in fair to excellent condition prior to the recent rainfall, although they were quite behind their normal stages of development. For those producers who are still able to get onto their land, in-crop spraying continues as weather and crop staging allow. Other sources of crop damage this week include hail and wind. Haying is just beginning for some producers and the majority of hay is rated as fair to good in quality.

Farmers are busy cleaning up flooded yards and homes, controlling weeds and getting ready for haying.

West-Central Saskatchewan (Crop Districts 6B – Hanley, Outlook, Loreburn, Saskatoon and Arelee areas; Crop District 7A – Rosetown, Kindersley, Eston, Major; CD 7B - Kerrobert, Macklin, Wilkie and Biggar areas)

Less than one per cent of the hay crop has been cut with 82 per cent rated as good in quality. While welcomed by some producers, recent rain has further delayed in-crop spraying. There are reports that aerial herbicide applications are increasing in some areas.

WC SK Crop Development - June 30			
	% Ahead	% Normal	% Behind
Fall Cereals	0	50	50
Spring Cereals	0	44	56
Oilseeds	0	43	57
Pulse Crops	2	53	45

Rainfall ranged from small amounts to 86 mm in the Smiley area. Since April 1, the Sonningdale area has received 282 mm of rain, the greatest amount for the region. Cropland topsoil moisture is rated as 12 per cent surplus, 87 per cent adequate and one per cent short. Hay land and pasture topsoil moisture is rated as seven per cent surplus and 93 per cent adequate.

The majority of crops are either at or behind their normal stages of development and are in fair to excellent condition. Warm and dry weather is needed to help advance

crops and for fieldwork to continue. Localized flooding, insects such as cutworms and diseases such as leaf spots and root rot have caused the majority of crop damage this week. Some producers are beginning to apply fungicides in earlier seeded crops while others are topdressing crops with fertilizer.

Farmers are busy controlling pests and beginning haying operations.

Northeastern Saskatchewan (Crop District 8 – Hudson Bay, Tisdale, Melfort, Carrot River, Humboldt, Kinistino, Cudworth and Aberdeen areas; Crop District 9AE – Prince Albert, Choiceland and Paddockwood areas)

Livestock producers in the northeast are just beginning haying operations in some areas. At this time, the majority of hay is rated as fair to good in quality. Recent rainfall has flooded many areas and there are reports of washed out roads and yards. There is water standing in many fields and crops are beginning to yellow due to excess moisture stress. The Star City area received 91 mm of rainfall over the weekend, bringing its total amount received since April 1 to 306 mm.

NE SK Crop Development - June 30			
	% Ahead	% Normal	% Behind
Fall Cereals	1	54	45
Spring Cereals	4	39	57
Oilseeds	3	38	59
Pulse Crops	0	50	50

Topsoil moisture conditions on cropland are worsening and are now rated as 80 per cent surplus and 20 per cent adequate. Hay land and pasture topsoil moisture is rated as 86 per cent surplus and 14 per cent adequate. CD 9AE is reporting that 93 per cent of the cropland and 96 per cent of the hay land and pasture has surplus topsoil moisture at this time. Low spots are covered in excess water and fields are very saturated.

Most crops are in poor to good condition and are behind their normal developmental stages. Crop damage this week is attributed to localized flooding and insects such as cutworms. In-crop spraying for both weeds and diseases may continue once fields are able to hold equipment.

Farmers are busy controlling pests, hauling grain and getting ready for haying.

Northwestern Saskatchewan (Crop District 9AW – Shellbrook, North Battleford, Big River and Hafford areas; Crop District 9B – Meadow Lake, Turtleford, Pierceland, Maidstone and Lloydminster areas)

Like much of the province, the northwestern region has received significant rainfall that delayed fieldwork for many producers. Haying operations have not begun but the majority of hay is rated as good to excellent in quality. Warm weather is needed to hasten crop, hay land and pasture maturity.

NW SK Crop Development - June 30			
	% Ahead	% Normal	% Behind
Fall Cereals	0	100	0
Spring Cereals	0	37	63
Oilseeds	0	35	65
Pulse Crops	0	41	59

The Turtleford area received 39 mm of rain this week while the Hafford area continues to lead the region in greatest total rainfall received since April 1 (336 mm). Cropland topsoil moisture conditions are improving and are now rated as 15 per cent surplus and 85 per cent adequate. Hay land and pasture topsoil moisture conditions are rated as five per cent surplus and 95 per cent adequate.

Crops continue to be delayed and most are either at or behind their normal developmental stages. Overall, crops are in good condition. There are reports of grasshoppers in the area, although damage is minimal at this time. Other sources of crop damage include localized flooding, wind and disease.

Farmers are busy controlling pests, getting ready for haying and hauling grain.

SK (Provincial) Crop Conditions - June 30, 2014							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	9	4	12	15	8	14	15
% good	47	46	57	61	55	54	36
% fair	28	46	22	19	28	23	26
% poor	14	4	7	4	8	8	17
% very poor	2	0	2	1	1	1	6
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	7	11	15	0	14	12	6
% good	50	51	71	51	60	54	75
% fair	26	27	14	29	17	20	19
% poor	10	8	0	13	6	6	0
% very poor	7	3	0	7	3	8	0

Southeast							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	13	13	4	11	1	6	16
% good	49	69	56	44	37	49	6
% fair	20	18	25	29	35	23	21
% poor	18	0	12	10	25	19	39
% very poor	0	0	3	6	2	3	18
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	9	7	52	0	3	1	0
% good	45	51	43	43	42	21	70
% fair	18	28	5	33	29	30	30
% poor	13	12	0	15	14	14	0
% very poor	15	2	0	9	12	34	0

Southwest							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	1	1	15	15	21	31	30
% good	68	40	69	71	74	55	41
% fair	28	59	14	12	5	11	24
% poor	3	0	2	2	0	3	5
% very poor	0	0	0	0	0	0	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	3	8	8	0	11	15	6
% good	74	80	77	95	71	64	75
% fair	21	11	15	5	15	17	19
% poor	2	1	0	0	3	4	0
% very poor	0	0	0	0	0	0	0

East-central							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	9	7	3	6	3	4	28
% good	30	17	46	40	46	46	50
% fair	37	51	37	37	38	39	18
% poor	17	25	13	17	12	10	4
% very poor	7	0	1	0	1	1	0
	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	1	3	12	0	17	9	0
% good	35	43	75	100	41	48	85
% fair	49	39	13	0	26	26	15
% poor	13	13	0	0	11	17	0
% very poor	2	2	0	0	5	0	0

West-central

	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	13	7	21	20	21	20	12
% good	79	73	71	62	70	70	52
% fair	8	20	8	18	9	10	31
% poor	0	0	0	0	0	0	5
% very poor	0	0	0	0	0	0	0

	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	24	22	19	10	22	18	20
% good	63	62	64	85	68	68	70
% fair	13	15	17	5	9	14	10
% poor	0	1	0	0	1	0	0
% very poor	0	0	0	0	0	0	0

Northeast

	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	0	0	2	0	2	1	0
% good	25	25	55	35	53	52	51
% fair	50	50	28	45	35	34	41
% poor	25	25	10	20	7	11	6
% very poor	0	0	5	0	3	2	2

	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	2	9	0	0	1	0	NA
% good	63	41	35	40	43	34	NA
% fair	22	34	30	35	43	30	NA
% poor	10	9	20	15	12	23	NA
% very poor	3	7	15	10	1	13	NA

Northwest							
	Winter wheat	Fall rye	Spring wheat	Durum	Oat	Barley	Canaryseed
% excellent	0	0	32	NA	26	33	NA
% good	50	100	53	NA	72	61	NA
% fair	50	0	11	NA	2	6	NA
% poor	0	0	3	NA	0	0	NA
% very poor	0	0	1	NA	0	0	NA

	Flax	Canola	Mustard	Soybean	Pea	Lentil	Chickpea
% excellent	30	29	NA	NA	25	NA	NA
% good	70	55	NA	NA	55	NA	NA
% fair	0	10	NA	NA	8	NA	NA
% poor	0	4	NA	NA	9	NA	NA
% very poor	0	2	NA	NA	3	NA	NA

Crop Districts and Rural Municipalities in Saskatchewan



Weekly Rainfall

for the week ending June 30, 2014



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Government of Saskatchewan

0 25 50 100 150 200

Kilometers

Projection: UTM Zone 13 Datum: NAD83

N

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Geomatics Services, Ministry of Agriculture July 2, 2014

© 2014 Government of Saskatchewan

Weekly Rainfall Summary

(in millimeters) 1 inch = 25 mm for the period June 24 to 30, 2014

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
1A	2	Mount Pleasant	97	226
	3	Enniskillen	52	177
	33	Moose Creek	75	244
	34	Browning	65	240
	61	Antler	195	430
	63	Moose Mountain	119	325
	64	Brock	82	242
	65	Tecumseh	96	314
1B	91	Maryfield	N/A	215
	122	Martin	260	520
	123	Silverwood	131	329
	124	Kingsley	140	320
	125 A	Chester	102	316
	125 B	Chester	100	342
	151 A	Rocanville	155	394
	151 B	Rocanville	134	434
	154	Elcapo	180	365
	155 A	Wolseley	135	304
	155 B	Wolseley	N/A	198
	67	Weyburn	N/A	196
2A	68	Brokenshell	11	198
	97	Wellington	55	214.5
2B	127 A	Francis	74	312.5
	127 B	Francis	51.5	215.4
	129	Bratt's Lake	38	234.5
	131 A	Baildon	50	282
	131 B	Baildon	40	297.2
	156 A	Indian Head	88.5	365
	156 B	Indian Head	132	403.5
	157	South Qu'Appelle	79	310
	160 A	Pense	98	299
	160 B	Pense	N/A	75
	161	Moose Jaw	N/A	365
	162	Caron	19	321
	191	Marquis	70	318
3ASE	38 A	Laurier	10	250.9
	38 B	Laurier	15	234
	39 A	The Gap	N/A	162
	39 B	The Gap	6	210
3ASW	10	Happy Valley	N/A	128
	12	Poplar Valley	51	253
	40 A	Bengough	16	184.5
	40 B	Bengough	25	224
	42	Willow Bunch	33	226
	43	Old Post	52	254
	70	Key West	N/A	32.4
	73 A	Stonehenge	35.3	240
	73 B	Stonehenge	34	306
	74	Wood River	N/A	111.5
3AN	101	Terrell	54	143
	102	Lake Johnston	N/A	141.1
	103	Sutton	N/A	131
	132 A	Hillsborough	19.5	216.5
	132 B	Hillsborough	33	281
	134	Shamrock	37	222
	193 A	Eyebrow	24	299
	193 B	Eyebrow	39	293
	224	Maple Bush	N/A	274
3BS	17	Val Marie	38	366
	75 A	Pinto Creek	17	246
	75 B	Pinto Creek	N/A	92
	76	Auvergne	15	180
	77	Wise Creek	N/A	208
	78	Grassy Creek	51	240
	105	Glenbain	9	178
	106	Whiska Creek	15	300
	107	Lac Pelletier	21	170
	108	Bone Creek	N/A	179
3BN	137	Swift Current	N/A	41.6
	138 A	Webb	N/A	173
	138 B	Webb	22	212.8
	166	Excelsior	46	195
	167	Sask. Landing	57	207
	168 A	Riverside	N/A	160
	168 B	Riverside	19.87	157.3
	226	Victory	N/A	206
	228	Lacadena	N/A	153
	257	Monet	20.25	123.1

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
4A	49	White Valley	16	142
	51	Reno	0.3	90.2
	79	Arlington	35	234
	109 A	Carmichael	N/A	177
	109 B	Carmichael	N/A	95
	110	Piapot	30	124
	111	Maple Creek	20	78
4B	139	Gull Lake	N/A	160
	141	Big Stick	24	133
	142	Enterprise	26	182
	169	Pittville	N/A	47
	231	Happyland	N/A	115
5A	183 A	Fertile Belt	139	332
	183 B	Fertile Belt	154	359
	186	Abernethy	117	357
	211	Churchbridge	87	294
	213	Saltcoats	167	374
	216	Tullymet	N/A	147.5
	241	Calder	79	258
	243	Wallace	148	311
	244	Orkney	N/A	239
	245 A	Garry	121	393
	245 B	Garry	131	416
	245 C	Garry	141	419
	246	Ituna Bon Accord	121	351
	247	Kellross	106	311
	248	Touchwood	89	271
5B	271	Cote	68	270
	273	Sliding Hills	191	477
	277	Emerald	154	497
	306	Invermay	96	384
	307	Elfros	82	358
	308 A	Big Quill	93	282
	308 B	Big Quill	104	265
	331	Livingston	N/A	236
	334	Preeceville	N/A	112
	336	Sasman	81	281
	337	Lakeview	N/A	199
	338	Lakeside	84	275
	366	Kelvington	87	298
	367	Ponass Lake	85	279
6A	190 A	Dufferin	154	460
	190 B	Dufferin	171	377.5
	190 C	Dufferin	99	335
	190 D	Dufferin	46	186.5
	219	Longlaketon	69	300
	220	Mckillop	112	433
	221	Sarnia	99.9	374.8
	222	Craik	72	320
	251	Big Arm	91	408
	252	Arm River	71	369
	279	Mount Hope	84	305
	282	McCraney	88	357
	309	Prairie Rose	83	350
	310	Usborne	N/A	154.5
	312	Morris	N/A	103
	313	Lost River	101	306
	339	Leroy	66.6	366.7
	340	Wolverine	41	293
	341	Viscount	42	268
	343 A	Blucher	29	262.9
	343 B	Blucher	30	140
6B	254	Lorebum	31	246
	284	Rudy	17	281
	285	Fertile Valley	16	197
	286	Milden	N/A	139
	314	Dundurn	22	226
	344	Corman Park	20	258
	346	Perdue	12	226
	376	Eagle Creek	24	265
	403	Rosthern	13	209

Crop Dist.	R.M. No.	Name	Past Week	Since 1-Apr
7A	259	Snipe Lake	N/A	20
	287	St. Andrews	20	260
	288	Pleasant Valley	74	222
	290 A	Kindersley	52	144
	290 B	Kindersley	55	105.5
	290 C	Kindersley	50	137.2
	292	Milton	70	185
	317 A	Marriott	N/A	78
	317 B	Marriott	12	170
	318	Mountain View	71	225
	320 A	Oakdale	N/A	106.1
	320 B	Oakdale	45	140
	321	Prairiedale	86	197
7B	347	Biggar	41	230
	350 A	Mariposa	17	110.6
	350 B	Mariposa	20	186
	351	Progress	49	157
	352	Heart's Hill	51	155
	377	Glenside	19	282
	378 A	Rosemount	26	172
	378 B	Rosemount	38	242
	379	Reford	22	222
	381	Grass Lake	N/A	103
	382	Eye Hill	45	164.1
	409	Buffalo	8	174
	410	Round Valley	38	225.4
8A	395	Porcupine	89	223
	397	Barrier Valley	68.4	282.9
	428	Star City	91	306
	456	Arborfield	72	276
	457	Connaught	62	209
	486	Moose Range	N/A	142
	487	Nipawin	55	193
8B	369	St. Peter	46	273
	370 A	Humboldt	41	299
	370 B	Humboldt	40	298
	371	Bayne	39	305
	372	Grant	65.5	297.5
	400	Three Lakes	29	271
	402	Fish Creek	56	290
	429	Flett's Springs	27	250
	459	Kinistino	40	288
	460	Birch Hills	43.2	241.9
9AE	488	Torch River	38	225
	491	Buckland	45	296
9AW	406	Mayfield	20	224
	435	Redberry	9	336
	436	Douglas	8	266
	463	Duck Lake	31	270.9
	467 A	Round Hill	N/A	316
	467 B	Round Hill	4	165
9B	438	Battle River	N/A	153
	440	Hillsdale	28	214
	442	Manitou Lake	12.3	158.4
	498 A	Parkdale	6	208.5
	498 B	Parkdale	7	167
	499	Mervin	2.5	142.5
	501 A	Frenchman Butte	N/A	216
	501 B	Frenchman Butte	39	176
	501 C	Frenchman Butte	6	213
	502	Britannia	5	147
	561	Loon Lake	N/A	214
	588 A	Meadow Lake	10	221
	588 B	Meadow Lake	N/A	232
	588 C	Meadow Lake	6	206
	588 D	Meadow Lake	19	273
	622	Beaver River	NIL	191.2

Municipality No: A, B, C and D - more than one reporter

These precipitation amounts represent point locations within each municipality and do not necessarily reflect the whole R. M.

Cumulative Rainfall

From: April 1, 2014
To: June 30, 2014



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Government of Saskatchewan

0 25 50 100 150 200
Kilometers
Projection: UTM Zone 13 Datum: NAD83

N

Data Source:
Rainfall - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Geomatics Services, Ministry of Agriculture July 2, 2014

© 2014 Government of Saskatchewan

Cropland Topsoil Moisture Conditions

July 1, 2014



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Government of Saskatchewan

0 25 50 100 150 200

Kilometers

Projection: UTM Zone 13 Datum: NAD83

N

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Geomatics Services, Ministry of Agriculture July 2, 2014

© 2014 Government of Saskatchewan

Hay and Pasture Topsoil Moisture Conditions

July 1, 2014



NOTE: Since techniques used to smooth the transition between zones can affect the values in localized areas, this map should be used for regional analysis only.

Government of Saskatchewan

0 25 50 100 150 200

Kilometers

Projection: UTM Zone 13 Datum: NAD83

N

Data Source:
Moisture - Ministry of Agriculture, Crop Report Database
IDW interpolation (power 2.5, fixed radius 300 km)

Geomatics Services, Ministry of Agriculture July 2, 2014

© 2014 Government of Saskatchewan